

Useful Nutrient Management Planning Data

Nutrient management plans should be based on accurate information. Ideally, manure generation and content should be determined on individual farms. Nutrients removed by harvesting crops also vary from farm to farm. The data provided here are general guidelines based on current research and can be used in preliminary planning.

Weights of crops (per bushel)

Crop	lb/bu	Crop	lb/bu
Corn	56	Rye	56
Sorghum	56	Sudangrass	40
Soybeans	60	Potatoes	60
Wheat	60	Sweetpotatoes	55
Barley	48	Sunflowers	25
Oats	32		

Length/Weight Conversion Factors

Length

- 1 inch = 2.54 centimeters
- 1 yard = 0.915 meters
- 1 mile = 5,280 feet = 1,610.7 meters
= 1.61 kilometers
- 1 meter = 100 centimeters = 1,000 millimeters
= 0.001 kilometers
- 1 chain = 66 feet = 100 links = 20.1 meters = 4 rods

Weight

- 1 pound = 454 grams = 0.454 kilograms = 16 ounces
- 1 ton (short) = 2,000 pounds
- 1 ton (long) = 2,240 pounds

More Speed Conversion Factors

- 1 mile per hour = 1.467 feet per second
= 88 feet per minute
= 26.8 meters per hour

Other

- 1 pound per acre = 1.12 kilograms per hectare
- 1 pound per gallon = 0.119 kilograms per liter
- parts per million (ppm) = micrograms per gram
= milligrams per liter
- 1 gallon per acre = 9.35 liter per hectare

Common Fertilizer Analyses

Fertilizer	Analysis	Chemical Formula
N		
Anhydrous ammonia	82-0-0	NH ₃
Ammonium nitrate	34-0-0	NH ₄ NO ₃
Urea	46-0-0	(NH ₂) ₂ CO
UAN solution (urea ammonium nitrate)	28 to 32-0-0	NH ₄ NO ₃ + (NH ₂) ₂ CO in water
Aqua ammonia	20-0-0	NH ₃ in water
Ammonium sulfate	21-0-0-24(S)	(NH ₄) ₂ SO ₄
P		
Triple superphosphate (TSP)	0-44 to 46-0	Ca(H ₂ PO ₄) ₂
Diammonium phosphate (DAP)	18-46-0	(NH ₄) ₂ HPO ₄
Monoammonium phosphate (MAP)	11-48-0	NH ₄ H ₂ PO ₄
Ammonium polyphosphate liquid (APP)	10-34-0	NH ₄ H ₂ PO ₄ + (NH ₄) ₃ HP ₂ O ₇
Ammonium polyphosphate dry (APP)	15-62-0	Same as liquid
K		
Potassium chloride (muriate of potash)	0-0-60	KCl
Potassium sulfate	0-0-50-18(S)	K ₂ SO ₄
Potassium-magnesium sulfate (sul-fo-mag)	0-0-22-22(S)-11(Mg)	K ₂ SO ₄ • 2MgSO ₄
Potassium nitrate	13-0-44	KNO ₃

Soil Testing Conversions

- Plow layer (6–7 inches) = parts per million X 2 = pounds per acre
- Top 12 inches = parts per million X 4 = pounds per acre



MISSISSIPPI STATE
UNIVERSITY
EXTENSION SERVICE

Manure Generation and Nutrient Content

Animal	Average Weight of Animal (lb)	Manure Production		Nutrient Content		
		(lb/d/1,000#)	(lb/d/1,000#)	N	P ₂ O ₅	K ₂ O
Dairy cow	1,200	80.00	1.30	0.45	0.16	0.31
Feeder swine	135	63.40	1.00	0.42	0.36	0.26
Gestation sow	375	27.20	0.44	0.19	0.14	0.14
Lactation sow w/ pigs	475	60.00	0.96	0.47	0.34	0.36
Nursery pig	20	106.00	1.70	0.60	0.57	0.42
Pullet litter	3	10.10	0.29	0.16	0.20	0.18
Breeder litter	8	13.40	0.38	0.21	0.27	0.23
Broiler litter	2	17.80	0.57	0.52	0.43	0.33

Nutrient Management Data

Plant Nutrient Removal (lb per acre unit of indicated yield)

	N	P ₂ O ₅	P	K ₂ O	K
Corn, 100 bushels grain	90	44	19	27	22
1 ton dry stover	22	8	4	32	26
Silage, ton	10	3.1	1	7	6
Cotton, bale	32	14	6	19	16
Rice, 150 bushels	86	45	20	24	20
Bahiagrass, ton	43	12	5	35	29
Hybrid bermudagrass	50	12	5	43	36
Sweetpotatoes, 100 bushels	24	13	6	56	46
Tall fescue	27	12	5	54	45
Soybeans, 40 bushels	152	34	15	52	43
Grain sorghum, 80 bushels	53	31	14	22	18
Wheat, 50 bushels	65	30	13	17	14
Peanuts, 6 tons	420	66	29	102	85
Dalligrass (5.8 tons/acre)	209	70	30	293	244
Annual ryegrass (4.5 tons/acre)	270	72	32	225	187

For more information, see MSU Extension Publication 2647 *Nutrient Management Guidelines for Agronomic Crops Grown in Mississippi*.

Fertilizer Conversion Factors

$$P_2O_5 \times 0.44 = P$$

$$P \times 2.29 = P_2O_5$$

$$K_2O \times 0.83 = K$$

$$K \times 1.20 = K_2O$$

$$1 \text{ gallon of water} = 8.328 \text{ pounds}$$

$$1 \text{ gallon of UAN (28\%N)} = 10.6 \text{ pounds}$$

Area/Volume Conversion Factors

Area

$$1 \text{ acre} = 43,560 \text{ feet}^2 = 0.405 \text{ hectare}$$

$$1 \text{ hectare} = 10,000 \text{ meters}^2$$

$$1 \text{ yard}^2 = 0.836 \text{ meter}^2$$

$$1 \text{ chain}^2 = 0.10 \text{ acre} = 16 \text{ rods}^2$$

$$1 \text{ mile}^2 = 640 \text{ acres}$$

Volume

$$1 \text{ bushel (level)} = 1.244 \text{ feet}^3 = 8 \text{ gallons (dry)}$$

$$= 9.31 \text{ gallons (liquid)} = 35.24 \text{ liters}$$

$$1 \text{ liter} = 1,000 \text{ milliliters or centimeters}^3$$

$$1 \text{ gallon (liquid)} = 3.78 \text{ liters} = 128 \text{ fluid ounces}$$

$$= 4 \text{ quarts} = 8 \text{ pints}$$

$$1 \text{ acre-foot} = 43,560 \text{ feet}^3 = 1,613 \text{ yards}^3$$

$$= 325,851 \text{ gallons}$$

$$1 \text{ cup} = 236.6 \text{ centimeters}^3 = 0.236 \text{ liters}$$

$$= 8 \text{ ounces} = 16 \text{ tablespoons}$$

Revised by **Larry Oldham**, PhD, Extension Soil Specialist, Department of Plant and Soil Sciences.

MAFES Special Research Initiatives and the Southwest Mississippi Research and Conservation District (through EPA Clean Water Act Section 319) provided grant funding for the project "Poultry Production By-Products Environmental Management Education Program Material Development."

Discrimination based upon race, color, religion, sex, national origin, age, disability, or veteran's status is a violation of federal and state law and MSU policy and will not be tolerated. Discrimination based upon sexual orientation or group affiliation is a violation of MSU policy and will not be tolerated.

Information Sheet 1620

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. GARY B. JACKSON, Director

(POD-04-14)



MISSISSIPPI STATE
UNIVERSITY

EXTENSION SERVICE

msucare.com